

## MEDICATION GUIDE

### Fenofibric Acid Delayed-Release Capsules, 45 mg and 135 mg Rx only

Read this Medication Guide before you start taking fenofibric acid delayed-release capsules and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment.

#### What is the most important information I should know about Fenofibric Acid Delayed-Release Capsules?

Fenofibric acid delayed-release capsules can be used with other cholesterol-lowering medicines called statins. Statins include:

- atorvastatin (Lipitor\*, Caduet\*)
- fluvastatin (Lescol\*, Lescol\* XL)
- lovastatin (Altoprev\*, Mevacor\*, Advicor\*)
- pitavastatin (Livalo\*)
- pravastatin (Pravachol\*)
- rosuvastatin (Crestor\*)
- simvastatin (Zocor\*, Simcor\*, Vytorin\*)

Statins can cause muscle pain, tenderness or weakness, which may be symptoms of a rare but serious muscle condition called rhabdomyolysis. In some cases rhabdomyolysis can cause kidney damage and death. The risk of rhabdomyolysis may be higher when fenofibric acid delayed-release capsules are given with statins. If you take a statin, tell your healthcare provider.

Other medicines or large amounts of grapefruit juice (more than a quart) may raise the levels of statins in your body, and could then raise the risk of muscle problems. Tell your healthcare provider if you are taking any medicines listed below.

- Heart medicine
- Stomach medicine
- Antibiotic
- Anti-fungal
- Cholesterol-lowering medicine
- Hormones
- HIV/AIDS medicine
- Antidepressant
- Immunosuppressant
- Anti-seizure medicine

Ask your healthcare provider or pharmacist for a list of these medicines, if you are not sure.

Tell your healthcare provider if you drink grapefruit juice.

#### What are Fenofibric Acid Delayed-Release Capsules?

Fenofibric acid delayed-release capsules are a prescription medicine used to treat cholesterol in the blood by lowering the total amount of triglycerides and LDL (bad) cholesterol, and increasing the HDL (good) cholesterol. **Fenofibric acid delayed-release capsules have not been shown to lower your risk of having heart problems or a stroke.** You should be on a low fat and low cholesterol diet while you take fenofibric acid delayed-release capsules.

The safety and effectiveness of fenofibric acid delayed-release capsules in children is not known.

#### Who should not take Fenofibric Acid Delayed-Release Capsules?

##### Do not take fenofibric acid delayed-release capsules if you:

- are allergic to fenofibric acid, or any of the ingredients in fenofibric acid delayed-release capsules. See the end of this Medication Guide for a list of all the ingredients in fenofibric acid delayed-release capsules.
- have severe kidney disease.
- have liver disease.
- have gallbladder disease.
- are a nursing mother.

##### Talk to your healthcare provider before you take fenofibric acid delayed-release capsules if you have any of these conditions.

##### What should I tell my healthcare provider before taking Fenofibric Acid Delayed-Release Capsules?

##### Before taking fenofibric acid delayed-release capsules, tell your healthcare provider about all your medical conditions, including if you:

- are allergic to any medicines.
- have ever had kidney problems.
- have ever had liver problems.
- have ever had gallbladder problems.
- are pregnant or if you plan to become pregnant. It is not known if fenofibric acid delayed-release capsules will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if fenofibric acid passes into your breast milk. You and your healthcare provider should decide if you will take fenofibric acid delayed-release capsules or breastfeed. You should not do both.

##### Tell your healthcare provider about all the medicines you take, including prescription and non-prescription medicines, vitamins and herbal supplements.

Using fenofibric acid delayed-release capsules with certain other medicines can affect the way these medicines work and other medicines may affect how fenofibric acid delayed-release capsules works. In some cases, using fenofibric acid delayed-release capsules with other medicines can cause serious side effects.

Know all the medicines you take. Keep a list of them and show it to your healthcare provider when you get a new medicine.

##### It is especially important to tell your healthcare provider if you take any of the medicines mentioned in, "What is the most important information I should know about Fenofibric Acid Delayed-Release Capsules?" or any of the medicines listed below:

- **anticoagulants**, also known as blood thinners (Warfarin, Coumadin)
- **bile acid resins**
- **cyclosporine**

Ask your healthcare provider if you are not sure if your medicine is one of these.

**TEMPLE Packaging Pvt. Ltd.** 28476001 Email:-[temple@bom4.vsnl.net.in](mailto:temple@bom4.vsnl.net.in)

Customer: Lupin Limited

Location: Goa

Product Code & Name: 222857 Fenofibric Acid DR Capsules MG(Booklet-Front)

SAP Code: 60100000604

Customer Artwork No.: xxxxx

File No.: 29337

Version No.: 13

Date: 02.05.2013

Open Size: 190 x 245 mm

No. of pages per Booklet : 24

Cover page printing : No

Barcode value: xxxx

Colours: 1 (Black)

Pharmacode value: xxxx

Substrate: 60 gsm Maplitho

Font: 10 pt helvetica condensed

Artwork Sr. No.: 141/9/12

Perforation: Yes (with printed line)

Artwork Status: final

Gluing: No

Note - This approval will be considered for final printing.

Please recheck for corrections indicated earlier, in this proof also.

Prepared by :

Q.C. Checked:

Checked by:

Approved by:

*shaji*

Customer Approval

Checked by:

Approved by:

Packaging Dvmp.t.

Production

Regulatory Affairs

Quality Assurance

**How should I take Fenofibric Acid Delayed-Release Capsules?**

- You should be on a low fat and low cholesterol diet while you take fenofibric acid delayed-release capsules.
- Take fenofibric acid delayed-release capsules one time each day as prescribed by your healthcare provider.
- Take fenofibric acid delayed-release capsules with or without food.
- Swallow fenofibric acid delayed-release capsules whole. Do not break, crush, dissolve, or chew fenofibric acid delayed-release capsules before swallowing. If you cannot swallow fenofibric acid delayed-release capsules whole, tell your healthcare provider, you may need a different medicine.
- If you take a medicine called a statin, you can take fenofibric acid delayed-release capsules and your statin at the same time of day.
- If you miss a dose of fenofibric acid delayed-release capsules, take it as soon as you remember. If it is almost time for your next dose, just skip the missed dose. Take the next dose at your regular time. If you are not sure about your dosing, call your healthcare provider. **Do not take more than one dose of fenofibric acid delayed-release capsules a day unless your healthcare provider tells you to.**
- If you take too much fenofibric acid delayed-release capsules, contact your healthcare provider or your local emergency department.
- Do not change your dose or stop fenofibric acid delayed-release capsules unless your healthcare provider tells you to.
- Your healthcare provider may do blood tests before you start taking fenofibric acid delayed-release capsules and during treatment. See your healthcare provider regularly to check your cholesterol and triglyceride levels and to check for side effects.

**What are the possible side effects with Fenofibric Acid Delayed-Release Capsules?**

**Fenofibric acid delayed-release capsules may cause serious side effects, including:**

- **muscle pain, tenderness, or weakness.** See “What is the most important information that I should know about fenofibric acid delayed-release capsules?”
- **tiredness and fever**
- **abdominal pain, nausea, or vomiting.** These may be signs of inflammation (swelling) of the gallbladder or pancreas.

**Call your healthcare provider right away if you have any of these serious side effects.**

The most common side effects with fenofibric acid delayed-release capsules include:

- headache
- heartburn (indigestion)
- nausea
- muscle aches
- increases in muscle or liver enzymes that are measured by blood tests

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of fenofibric acid delayed-release capsules.

For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**How do I store Fenofibric Acid Delayed-Release Capsules?**

- Store fenofibric acid delayed-release capsules between 59 to 86° F (15 to 30° C).
- Protect fenofibric acid delayed-release capsules from moisture.

**Keep fenofibric acid delayed-release capsules and all medicines out of the reach of children.**

**General information about the safe and effective use of Fenofibric Acid Delayed-Release Capsules.**

Medicines are sometimes prescribed for conditions that are not mentioned in the Medication Guide. Do not use fenofibric acid delayed-release capsules for a condition for which it was not prescribed. Do not give fenofibric acid delayed-release capsules to other people, even if they have the same condition you have. It may harm them.

This Medication Guide summarizes the most important information about fenofibric acid delayed-release capsules. If you would like more information, talk to your healthcare provider. You can also ask your pharmacist or healthcare provider for information that is written for health professionals.

For more information go to [www.lupinpharmaceuticals.com](http://www.lupinpharmaceuticals.com) or call at 1-800-399-2561.

**What are the ingredients in Fenofibric Acid Delayed-Release Capsules?**

**Active Ingredient:** Fenofibric acid

**Inactive Ingredients:** colloidal silicon dioxide, dibutyl sebacate, ethyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, methacrylic acid co polymer, povidone, sodium stearyl fumarate, talc and triethyl citrate. The capsule shell of the 45 mg capsule contains the following inactive ingredients: gelatin, iron oxide black, iron oxide red, iron oxide yellow, sodium lauryl sulphate and titanium dioxide. The capsule shell of the 135 mg capsule contains the following inactive ingredients: FD and C Blue #1, gelatin, iron oxide yellow, sodium lauryl sulphate and titanium dioxide. The capsules are printed with edible ink containing iron oxide black, potassium hydroxide, propylene glycol and shellac.

\* The brands listed are trademarks of their respective owners and are not trademarks of Lupin Pharmaceuticals, Inc. The makers of these brands are not affiliated with and do not endorse Lupin Pharmaceuticals, Inc. or its products.

**Manufactured for  
Lupin Pharmaceuticals, Inc.**  
Baltimore, Maryland 21202  
United States

Manufactured by:  
**Lupin Limited**  
Goa - 403 722  
INDIA

May 2013

ID#: 222857

|  |                        |   |                   |
|--|------------------------|---|-------------------|
| <b>TEMPLE Packaging Pvt. Ltd.</b> 28476001 Email:- <a href="mailto:temple@bom4.vsnl.net.in">temple@bom4.vsnl.net.in</a>              |                        |   |                   |
| <b>Customer:</b> Lupin Limited   |                        | <b>Location:</b> Goa                        |                   |
| <b>Product Code &amp; Name:</b> 222857 Fenofibric Acid DR Capsules MG(Booklet-Back)  |                        |   |                   |
| <b>SAP Code:</b> 60100000604   |                        | <b>Customer Artwork No.:</b> xxxxx          |                   |
| <b>File No.:</b> 29337   | <b>Version No.:</b> 13 | <b>Date:</b> 02.05.2013                     |                   |
| <b>Open Size:</b> 190 x 245 mm   |                        | <b>No. of pages per Booklet :</b> 24        |                   |
| <b>Cover page printing :</b> No  |                        | <b>Barcode value:</b> xxxx                  |                   |
| <b>Colours:</b> 1 (Black)  |                        | <b>Pharmacode value:</b> xxxx               |                   |
| <b>Substrate:</b> 60 gsm Maplitho  |                        | <b>Font:</b> 10 pt helvetica condensed      |                   |
| <b>Artwork Sr. No.:</b> 141/9/12   |                        | <b>Perforation:</b> Yes (with printed line) |                   |
| <b>Artwork Status:</b> final   |                        | <b>Gluing:</b> No                           |                   |
| Note - This approval will be considered for final printing.<br>Please recheck for corrections indicated earlier, in this proof also. |                        |   |                   |
| Prepared by :  | Q.C. Checked:          | Checked by:                                 | Approved by:      |
| <i>shaji</i>   |                        |   |                   |
| Customer Approval  |                        |   |                   |
| Checked by:  |                        | Approved by:                                |                   |
| Packaging Dvmpnt.  | Production             | Regulatory Affairs                          | Quality Assurance |