

(Front of card)

**WARNING CARD**  
**Abacavir, Lamivudine and Zidovudine Tablets**  
**300 mg/150 mg/300 mg**

**Patients taking abacavir, lamivudine and zidovudine tablets may have a serious allergic reaction (hypersensitivity reaction) that can cause death. If you get a symptom from 2 or more of the following groups while taking abacavir, lamivudine and zidovudine tablets, call your healthcare provider right away to find out if you should stop taking this medicine.**

	Symptom(s)
<b>Group 1</b>	<b>Fever</b>
<b>Group 2</b>	<b>Rash</b>
<b>Group 3</b>	<b>Nausea, vomiting, diarrhea, or abdominal (stomach area) pain</b>
<b>Group 4</b>	<b>Generally ill feeling, extreme tiredness, or achiness</b>
<b>Group 5</b>	<b>Shortness of breath, cough, or sore throat</b>

Always carry this Warning Card with you to help recognize symptoms of this allergic reaction.

(Back of Card)

**WARNING CARD**  
**Abacavir, Lamivudine and Zidovudine Tablets**

If you must stop treatment with abacavir, lamivudine and zidovudine tablets because you have had an allergic reaction to abacavir, **NEVER** take abacavir, lamivudine and zidovudine tablets or any other abacavir-containing medicine (ZIAGEN<sup>®</sup>, EPZICOM<sup>®</sup>, or TRIUMEQ<sup>®</sup>) again. If you have an allergic reaction, dispose of any unused abacavir, lamivudine and zidovudine tablets. Ask your pharmacist how to properly dispose of medicines. If you take abacavir, lamivudine and zidovudine tablets or another abacavir-containing medicine again after you have had an allergic reaction, **WITHIN HOURS** you may get **life-threatening symptoms** that may include **very low blood pressure or death**.

Please read the Medication Guide for additional information on abacavir, lamivudine and zidovudine tablets.

The brands listed are trademarks of their respective owners and are not trademarks of Lupin Pharmaceuticals, Inc. The makers of these brands are not affiliated with and do not endorse Lupin Pharmaceuticals, Inc. or its products.

July 2017

ID#: 252302